

PC26084A

AMINOPYRAZOLE DERIVATIVES AS
GSK-3 INHIBITORS

5

FIELD OF THE INVENTION

The invention relates to certain 5-cyclobutyl-3-aminopyrazoles which inhibit kinases, such as glycogen synthase kinase-3 (GSK-3), cyclin-dependent kinase-2 (cdk-2), and cyclin-dependent kinase-5 (cdk-5). As such, the compounds are useful in the treatment of conditions, diseases, and symptoms including, *inter alia*,
10 Alzheimer's Disease, cancer, dementia, depression, diabetes, hair loss, schizophrenia, and stroke.


BACKGROUND OF THE INVENTION

The serine/threonine kinase cdk-2 is essential for normal cellular cycling and
15 plays a critical role in disorders arising from abnormal cell cycling, a common characteristic of many oncological disorders. Inhibitors of cdk-2 are therefore useful in the treatment of various types of cancers and other diseases or conditions related to abnormal cell growth. See, for example, Meijer, et al., Pharmacol. and Therapeutics, 82 (2-3), 279-284 (1999), Sausville, et al., Pharmacol. and Therapeutics, 82 (2-3),
20 285-292 (1999). The serine/threonine kinase cdk-5, along with its cofactor p25, or the longer cofactor p35, has been linked to neurodegenerative disorders, and inhibitors of cdk-5 are therefore useful in the treatment of disorders such as Alzheimer's Disease, Parkinson's Disease, stroke, and Huntington's Disease. Treatment of such neurodegenerative disorders using cdk-5 inhibitors is supported by the finding that
25 cdk-5 is involved in the phosphorylation of tau protein, and dopamine and cyclic AMP-regulated phosphoprotein (DARPP-32) at threonine 75, and is thus indicated as playing a role in dopaminergic transmission.

Glycogen synthase kinase-3 (GSK-3), a proline-directed, serine/threonine kinase for which two isoforms, GSK-3 α and GSK-3 β , have been identified,
30 phosphorylates the rate-limiting enzyme of glycogen synthesis, glycogen synthase (GS). See, for example, Embi, et al., Eur. J. Biochem., 107, 519-527 (1980). GSK-3 α and GSK-3 β are both highly expressed in the body. See, for example, Woodgett, et al., EMBO, 9, 2431-2438 (1990) and Loy, et al., J. Peptide Res., 54, 85-91 (1999). Besides GS, a number of other GSK-3 substrates have been identified, including

many metabolic, signaling, and structural proteins. Notable among the plurality of signaling proteins regulated by GSK-3 are many transcription factors, including activator protein-1; cyclic AMP response element binding protein (CREB); the nuclear factor (NF) of activated T-cells; heat shock factor-1; β -catenin; c-Jun; c-Myc; c-Myb; and NF- κ B. See, for example, C. A. Grimes, et al., *Prog. Neurobiol.*, 65, 391-426 (2001), H. Eldar-Finkelman, *Trends in Molecular Medicine*, 8, 126-132 (2002), and P. Cohen, et al., *Nature*, 2, 1-8, (2001). Accordingly, targeting the activity of GSK-3 has significant therapeutic potential in the treatment of many disparate pathologies and conditions, for example, Alzheimer's Disease (A. Castro, et al., *Exp. Opin. Ther. Pat.*, 10, 1519-1527 (2000)); asthma (P. J. Barnes, *Ann. Rev. Pharmacol. Toxicol.*, 42, 81-98 (2002)); cancer (Beals, et al., *Science*, 275, 1930-1933 (1997), L. Kim, et al., *Curr. Opin. Genet. Dev.*, 10, 508-514 (2000), and Q. Eastman, et al., *Curr. Opin. Cell Biol.*, 11, 233 (1999)); diabetes and its related sequelae, for example, Syndrome X and obesity (S. E. Nikoulina, et al., *Diabetes*, 51, 2190-2198 (2002), Orena, et al., *JBC*, 15765-15772 (2000), and Summers, et al., *J. Biol. Chem.*, 274, 17934-17940 (1999)); hair loss (S. E. Millar, et al., *Dev. Biol.*, 207, 133-149 (1999) and E. Fuchs, et al., *Dev. Cell*, 1, 13-25 (2001)); inflammation (P. Cohen, *Eur. J. Biochem.*, 268, 5001-5010 (2001)); mood disorders, such as depression (A. Adnan, et al., *Chem. Rev.*, 101, 2527-2540 (2001) and R. S. B. Williams, et al., *Trends Pharmacol. Sci.*, 21, 61-64 (2000)); neuronal cell death and stroke (D. A. E. Cross, et al., *J. Neurochem.*, 77, 94-102 (2001) and C. Sasaki, et al., *Neurol. Res.*, 23, 588-592 (2001)); bipolar disorder (Klein, et al., *PNAS*, 93, 8455-8459 (1996)); skeletal muscle atrophy (G. J. Brunn, et al., *Science*, 277, 99-101 (1997), R. E. Rhoads, *J. Biol. Chem.*, 274, 30337-30340 (1999), V. R. Dharmesh, et al., *Am. J. Physiol. Cell Physiol.*, 283, C545-551 (2002), and K. Baar, et al., *A. J. Physiol.*, 276, C120-C127 (1999)); decreased sperm motility (Vijayaraghavan, et al., *Biol. Reproduction*, 54, 709-718 (1996)); and in cardio-protection (C. Badorff, et al., *J. Clin. Invest.*, 109, 373-381 (2002), S. Haq, et al., *J. Cell Biol.*, 151, 117-129 (2000), and H. Tong, et al., *Circulation Res.*, 90, 377-379 (2002)).

Certain pyrazole derivatives of formula (II),




useful as inhibitors of cdk2, cdk5, and GSK-3, are disclosed in commonly-assigned PCT International Application Publication No. WO 02/18346, wherein R¹, R², R³, and R⁴ are as defined therein.

5

SUMMARY OF THE INVENTION

The present invention provides compounds of formula (I)




10

the stereoisomers and prodrugs thereof, and the pharmaceutically acceptable salts of the compounds, stereoisomers, and prodrugs, wherein R¹, R², and R³ are as defined herein; pharmaceutical compositions thereof; combinations thereof; and uses thereof in the treatment of, *inter alia*, conditions, diseases, and symptoms including, *inter alia*, Alzheimer's Disease, cancer, dementia, depression, diabetes, hair loss, schizophrenia, and stroke.

15

DETAILED DESCRIPTION OF THE INVENTION

The invention provides compounds of formula (I)



the stereoisomers and prodrugs thereof, and the pharmaceutically acceptable salts of the compounds, stereoisomers, and prodrugs, wherein:

20

R¹ is:

- (A) -(C₁-C₆)alkyl, optionally substituted independently with from one to three (a) halogen; (b) heteroaryl, optionally substituted independently with from one to three -(C₁-C₆)alkyl; trifluoromethyl; or -(C₁-C₆)alkoxy; (c) aryl, optionally substituted independently with from one to three halogen; -(C₁-C₆)alkoxy; trifluoromethyl; -(C₁-C₆)alkyl; or -C(O)(C₁-C₆)alkyl; (d) -OR⁵; (e) -(C₃-C₈)cycloalkyl; or (f) heterocycloalkyl;

25

(B) $-(C_3-C_8)$ cycloalkyl, optionally substituted independently with from one to three (g) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)$ alkyl; trifluoromethyl; or $-(C_1-C_6)$ alkoxy; (h) aryl, optionally substituted independently with from one to three halogen; $-(C_1-C_6)$ alkoxy; trifluoromethyl; $-(C_1-C_6)$ alkyl; or $-C(O)(C_1-C_6)$ alkyl; (i) heterocycloalkyl; (j) $-OR^5$; or (k) $-(C_1-C_6)$ alkyl, optionally substituted with from one to three halogen;

(C) heterocycloalkyl, optionally substituted with from one to three (l) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)$ alkyl; trifluoromethyl; or $-(C_1-C_6)$ alkoxy; (m) aryl, optionally substituted independently with from one to three halogen; $-(C_1-C_6)$ alkoxy; trifluoromethyl; $-(C_1-C_6)$ alkyl; or $-C(O)(C_1-C_6)$ alkyl; (n) $-(C_3-C_8)$ cycloalkyl; (o) heterocycloalkyl; (p) $-OR^5$; or (q) $-(C_1-C_6)$ alkyl, optionally substituted with from one to three halogen; or

(D) heteroaryl, optionally substituted with from one to three $-(C_1-C_6)$ alkyl or trifluoromethyl;

15 R^2 and R^3 are, independently,

(E) hydrogen;

(F) $-(C_1-C_6)$ alkyl, optionally substituted independently with from one to three (r) halogen; (s) aryl, optionally substituted independently with from one to three halogen; trifluoromethyl; $-(C_1-C_6)$ alkyl, or $-(C_1-C_6)$ alkoxy, optionally substituted with from one to three fluorine atoms; (t) heteroaryl, optionally substituted independently with from one to three nitro; $-(C_1-C_6)$ alkyl; trifluoromethyl; halogen; or $-(C_1-C_6)$ alkoxy; (u) heterocycloalkyl, optionally substituted independently with one to three $-(C_1-C_6)$ alkyl; oxo; aryl; or heteroaryl; (v) $-(C_3-C_8)$ cycloalkyl, optionally substituted independently with from one to three cyano or aryl; (w) $-NHR^4$; (x) $-OR^5$; (y) $-N[(C_1-C_6)alkyl]_2$; or (z) cyano;

(G) $-(C_3-C_8)$ cycloalkyl, optionally substituted independently with from one to three cyano or aryl;

(H) aryl, optionally substituted independently with from one to three halogen; $-(C_1-C_6)$ alkoxy; trifluoromethyl; or $-(C_1-C_6)$ alkyl;

30 (I) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)$ alkyl or $-(C_1-C_6)$ alkoxy; or

(J) heterocycloalkyl, optionally substituted with from one to three $-(C_1-C_6)$ alkyl, optionally substituted with aryl; or

R^2 and R^3 , taken together with the nitrogen atom to which they are attached, form a heterocycloalkyl ring, optionally substituted independently with (aa) $-(C_1-C_6)alkyl$, optionally substituted with $-R^4$ or $-OR^5$; (bb) aryl; (cc) heteroaryl; (dd) $-N[(C_1-C_6)alkyl]R^4$; (ee) $-R^4$; or (ff) $-(C_1-C_6)alkoxy$;

5 R^4 is (K) $-(C_1-C_6)alkyl$; (L) $-C(O)(C_1-C_6)alkyl$; (M) $-C(O)O(C_1-C_6)alkyl$, optionally substituted with aryl; (N) aryl; (O) heteroaryl; or (P) heterocycloalkyl, wherein each (N) aryl, (O) heteroaryl, or (P) heterocycloalkyl group is optionally substituted independently with from one to three (gg) halogen; (hh) nitro; (ii) trifluoromethyl; (jj) $-(C_1-C_6)alkyl$; or (kk) $-N[(C_1-C_6)alkyl][C(O)(C_1-C_6)alkyl]$; and

10 R^5 is (Q) $-(C_1-C_6)alkyl$; (R) $-C(O)(C_1-C_6)alkyl$; (S) aryl; (T) heteroaryl; or (U) heterocycloalkyl, wherein each (S) aryl, (T) heteroaryl, or (U) heterocycloalkyl group is optionally substituted independently with from one to three (ll) halogen; (mm) nitro; (nn) trifluoromethyl; (oo) $-(C_1-C_6)alkyl$; or (pp) $-N[(C_1-C_6)alkyl][C(O)(C_1-C_6)alkyl]$.

A generally preferred subgroup of the compounds of formula (I) comprises
15 those compounds wherein:

R^1 is:

(A) $-(C_1-C_6)alkyl$, optionally substituted independently with (b) heteroaryl, optionally substituted independently with $-(C_1-C_6)alkyl$; trifluoromethyl; or $-(C_1-C_6)alkoxy$; (c) aryl, optionally substituted independently with from one to three
20 halogen; $-(C_1-C_6)alkoxy$; trifluoromethyl; $-(C_1-C_6)alkyl$; (d) $-OR^5$; or (f) heterocycloalkyl;

(B) $-(C_3-C_8)cycloalkyl$, optionally substituted independently with (g) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)alkyl$; trifluoromethyl; or $-(C_1-C_6)alkoxy$; (h) aryl, optionally substituted independently with from one to three halogen; $-(C_1-C_6)alkoxy$; trifluoromethyl; or $-(C_1-C_6)alkyl$; (i)
25 heterocycloalkyl; (j) $-OR^5$; (k) $-(C_1-C_6)alkyl$, optionally substituted with from one to three halogen;

(C) heterocycloalkyl, optionally substituted with (l) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)alkyl$; trifluoromethyl; or $-(C_1-C_6)alkoxy$; (m) aryl, optionally substituted independently with from one to three
30 halogen; $-(C_1-C_6)alkoxy$; trifluoromethyl; $-(C_1-C_6)alkyl$; or $-C(O)(C_1-C_6)alkyl$; (n) $-(C_3-C_8)cycloalkyl$; (o) heterocycloalkyl; (p) $-OR^5$; or (q) $-(C_1-C_6)alkyl$, optionally substituted with from one to three halogen;

R^2 is hydrogen or $-(C_1-C_6)alkyl$;

R^3 is:

(F) $-(C_1-C_6)alkyl$, optionally substituted independently with from one to three (r) halogen; (s) aryl, optionally substituted independently with from one to three halogen; trifluoromethyl; $-(C_1-C_6)alkyl$, or $-(C_1-C_6)alkoxy$, optionally substituted with from one to three fluorine atoms; (t) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)alkyl$; trifluoromethyl; halogen; or $-(C_1-C_6)alkoxy$; (u) heterocycloalkyl, optionally substituted independently with one to three $-(C_1-C_6)alkyl$; oxo; aryl; or heteroaryl; (v) $-(C_3-C_8)cycloalkyl$; (w) $-NHR^4$; (x) $-OR^5$; (y) $-N[(C_1-C_6)alkyl]_2$; or (z) cyano;

(G) $-(C_3-C_8)cycloalkyl$, optionally substituted independently with from one to three cyano or aryl; or

(J) heterocycloalkyl, optionally substituted with from one to three $-(C_1-C_6)alkyl$, optionally substituted with aryl; or

R^2 and R^3 , taken together with the nitrogen atom to which they are attached, form a heterocycloalkyl ring, optionally substituted independently with (aa) $-(C_1-C_6)alkyl$, optionally substituted with $-R^4$ or $-OR^5$; (bb) aryl; (cc) heteroaryl; or (ff) $-(C_1-C_6)alkoxy$;

R^4 is (K) $-(C_1-C_6)alkyl$; (N) aryl; (O) heteroaryl; or (P) heterocycloalkyl, wherein each aryl, heteroaryl, or heterocycloalkyl group is optionally substituted independently with from one to three (gg) halogen; (ii) trifluoromethyl; or (jj) $-(C_1-C_6)alkyl$; and

R^5 is (Q) $-(C_1-C_6)alkyl$; (S) aryl; (T) heteroaryl; or (U) heterocycloalkyl, wherein each (S) aryl, (T) heteroaryl, or (U) heterocycloalkyl group is optionally substituted independently with from one to three (ll) halogen; (nn) trifluoromethyl; or (oo) $-(C_1-C_6)alkyl$.

Another generally preferred subgroup of the compounds of formula (I) comprises those compounds wherein:

R^1 is:

(A) $-(C_1-C_6)alkyl$, optionally substituted independently with (b) heteroaryl, optionally substituted independently with $-(C_1-C_6)alkyl$ or $-(C_1-C_6)alkoxy$; (c) aryl, optionally substituted independently with from one to three halogen; $-(C_1-C_6)alkoxy$; trifluoromethyl; or $-(C_1-C_6)alkyl$; or (d) $-OR^5$;

(B) $-(C_3-C_8)cycloalkyl$, optionally substituted independently with (g) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)alkyl$ or $-(C_1-C_6)alkoxy$; (h) aryl, optionally substituted independently with from one to three

halogen; $-(C_1-C_6)$ alkoxy; trifluoromethyl; or $-(C_1-C_6)$ alkyl; (j) $-OR^5$; (k) $-(C_1-C_6)$ alkyl, optionally substituted with from one to three halogen; or

(C) heterocycloalkyl, optionally substituted with (l) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)$ alkyl or $-(C_1-C_6)$ alkoxy; (m) aryl, optionally substituted independently with from one to three halogen; $-(C_1-C_6)$ alkoxy; trifluoromethyl; or $-(C_1-C_6)$ alkyl; (p) $-OR^5$; or (q) $-(C_1-C_6)$ alkyl, optionally substituted with from one to three halogen;

R^2 is hydrogen or $-(C_1-C_6)$ alkyl;

R^3 is:

(F) $-(C_1-C_6)$ alkyl, optionally substituted independently with (s) aryl, optionally substituted independently with from one to three halogen; trifluoromethyl; $-(C_1-C_6)$ alkyl, or $-(C_1-C_6)$ alkoxy, optionally substituted with from one to three fluorine atoms; (t) heteroaryl, optionally substituted independently with from one to three $-(C_1-C_6)$ alkyl or trifluoromethyl; and

R^5 is (S) aryl, optionally substituted with halogen.

The compounds and intermediates of the present invention may be named according to either the IUPAC (International Union for Pure and Applied Chemistry) or CAS (Chemical Abstracts Service, Columbus, OH) nomenclature systems.

The carbon atom content of the various hydrocarbon-containing moieties may be indicated by a prefix designating the minimum and maximum number of carbon atoms in the moiety, i.e., the prefix " $-(C_a-C_b)$ alkyl" indicates an alkyl moiety of the integer "a" to "b" carbon atoms, inclusive.

The term "alkoxy" denotes straight or branched, monovalent, saturated aliphatic chains of carbon atoms bonded to an oxygen atom, wherein the alkoxy group optionally incorporates one or more double or triple bonds, or a combination of double bonds and triple bonds. Examples of alkoxy groups include methoxy, ethoxy, propoxy, butoxy, *iso*-butoxy, *tert*-butoxy, and the like.

The term "alkyl" denotes straight, or branched, monovalent chains of carbon atoms, wherein the alkyl group optionally incorporates one or more double or triple bonds, or a combination of double bonds and triple bonds. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, vinyl, allyl, 2-methylpropenyl, 2-butenyl, 1,3-butadienyl, ethynyl, propargyl, and the like.

The term "aryl" denotes a monocyclic, or polycyclic, aromatic hydrocarbon. Examples of aryl groups include anthracenyl, fluorenyl, phenanthrenyl, phenyl, naphthyl, and the like.

5 The term "cycloalkyl" denotes a saturated monocyclic, or polycyclic, cycloalkyl group, optionally fused to an aryl group, wherein the cycloalkyl group optionally incorporates one or more double or triple bonds, or a combination of double bonds and triple bonds, but which is not aromatic. Examples of cycloalkyl groups include adamantanyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, decahydronaphthalinyl, norbornanyl, and the like.

10 The term "halogen" represents chloro, fluoro, bromo, and iodo.

The term "heteroaryl" denotes a monocyclic, or polycyclic, aromatic hydrocarbon group wherein one or more carbon atoms have been replaced with heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur. If the heteroaryl group contains more than one heteroatom, the heteroatoms may be the
15 same or different. Examples of heteroaryl groups include acridinyl, benzofuranyl, benzothienyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, chromenyl, cinnolinyl, furyl, imidazolyl, indazolyl, indoliziny, indolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthyridinyl, oxadiazolyl, oxazinyl, oxazolyl, phenazinyl, phthalazinyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrido[3,4-
20 b]indolyl, pyridyl, pyrimidyl, pyrrolyl, quinoliziny, quinolyl, quinoxaliny, thiadiazolyl, thiatriazolyl, thiazolyl, thienyl, triazinyl, triazolyl, xanthenyl, and the like.

The term "heterocycloalkyl" denotes a saturated, or partially unsaturated, monocyclic, or polycyclic, cycloalkyl group, optionally fused to an aromatic or heteroaromatic hydrocarbon group, in which at least one of the carbon atoms has
25 been replaced with a heteroatom selected from the group consisting of nitrogen, oxygen, and sulfur. If the heterocycloalkyl group contains more than one heteroatom, the heteroatoms may be the same or different. Examples of such heterocycloalkyl groups include azabicycloheptanyl, azetidiny, benzazepiny, 1,3-dihydroisoindolyl, dioxolanyl, dioxanyl, carbazolyl, dioxolanyl, dithianyl, indoliny, imidazolidiny, morpholiny, quinuclidiny, phenothiaziny, phenoxaziny, piperaziny, piperidyl,
30 pyrazolidiny, pyrrolidiny, tetrahydrofuryl, tetrahydroindolyl, tetrahydroisoquinoliny, tetrahydropyranly, tetrahydroquinoliny, tetrahydroquinoxaliny, tetrahydrothiopyranly, tetrahydro-2H-1,4-thiaziny, thiazolidiny, thiomorpholiny, thioxanthenyl, thioxanyl, trithianyl, and the like.

A cyclic group may be bonded to another group in more than one way. If no particular bonding arrangement is specified, then all possible arrangements are intended. For example, the term "pyridyl" includes 2-, 3-, or 4-pyridyl, and the term "thienyl" includes 2- or 3-thienyl.

5 The term "mammal" means animals including, for example, dogs, cats, cows, sheep, horses, and humans. Preferred mammals include humans of either gender.

 The term "oxo", when used within the context of the term "heterocycloalkyl", indicates a carbonyl substituent formed between a ring carbon atom(s) of the heterocycloalkyl group and an oxygen atom.

10 The phrase "pharmaceutically acceptable" indicates that the designated carrier, vehicle, diluent, excipient(s), and/or salt must be chemically and/or physically compatible with the other ingredients comprising the formulation, and physiologically compatible with the recipient thereof.

 The term "prodrug" refers to a compound that is a drug precursor which,
15 following administration, releases the drug *in vivo* via a chemical or physiological process (e.g., upon being brought to physiological pH or through enzyme activity). A discussion of the preparation and use of prodrugs is provided by T. Higuchi and W. Stella, "Prodrugs as Novel Delivery Systems, Vol. 14 of the ACS Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American
20 Pharmaceutical Association and Pergamon Press, 1987.

 The term "radical" denotes a group of atoms that behaves as a single atom in a chemical reaction, e. g., an organic radical is a group of atoms that imparts characteristic properties to a compound containing it, or which remains unchanged during a series of reactions, or transformations.

25 The term "salts" refers to organic and inorganic salts of a compound of formula (I), or a prodrug thereof. These salts can be prepared *in situ* during the final isolation and purification of a compound, or by separately reacting a compound of formula (I), or a prodrug thereof, with a suitable organic or inorganic acid or base and isolating the salt thus formed. Representative salts include the hydrobromide,
30 hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, besylate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts, and the like. These may also include cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium,

magnesium, and the like, as well as non-toxic ammonium, quaternary ammonium, and amine cations including, but not limited to, ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. For additional examples see, for example, Berge, et al., J. Pharm. Sci., 66, 1-19 (1977).

The term "substituted" means that a hydrogen atom on a molecule has been replaced with a different atom or molecule. The atom or molecule replacing the hydrogen atom is denoted as a "substituent."

The symbol "-" represents a covalent bond.

The phrase "reaction-inert solvent" or "inert solvent" refers to a solvent, or mixture of solvents, that does not interact with starting materials, reagents, intermediates, or products in a manner that adversely affects their desired properties.

The terms "treating", "treated", or "treatment" as employed herein includes preventative (e.g., prophylactic), palliative, or curative use or result.

The compounds of formula (I) may contain asymmetric or chiral centers and, therefore, exist in different stereoisomeric forms. It is intended that all stereoisomeric forms of the compounds and prodrugs of formula (I) as well as mixtures thereof, including racemic mixtures, form part of the present invention. In addition, the present invention embraces all geometric and positional isomers. For example, if a compound or prodrug of formula (I) incorporates a double bond, both the *cis*- and *trans*- forms, as well as mixtures thereof, are embraced within the scope of the invention.

Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well-known to those of ordinary skill in the art, such as by chromatography and/or fractional crystallization.

Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of formula (I) may be atropisomers (e.g., substituted biaryls) and are also considered as part of the invention.

The compounds and prodrugs of formula (I) may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents, such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms.

It is also possible that the compounds and prodrugs of formula (I) may exist as tautomeric isomers in equilibrium, and all such forms are embraced within the scope of the invention.

The present invention also embraces isotopically-labeled compounds of formula (I), which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of formula (I) include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine, and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. The compounds of formula (I), the prodrugs thereof, and the pharmaceutically acceptable salts of the compounds and prodrugs, that contain the aforementioned isotopes and/or other isotopes of the other atoms are intended to be within the scope of the instant invention.

Certain isotopically-labeled compounds of formula (I), for example those compounds into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in compound and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their relative ease of preparation and facile detection. Furthermore, substitution with heavier isotopes such as deuterium, i.e., ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life, or reduced dosage requirements and, hence, may be preferred in some circumstances. The isotopically-labeled compounds of formula (I) can generally be prepared by carrying out procedures analogous to those disclosed in the Schemes and/or Examples set forth hereinbelow, by substituting an isotopically-labeled reagent for a non-isotopically-labeled reagent.

In another aspect, the invention provides methods for inhibiting cdk2, cdk5, and/or GSK-3 activity in a mammal in need of such inhibition which methods comprise administering to the mammal a cdk2, cdk5, and/or GSK-3 activity inhibiting amount of a compound of formula (I), a prodrug thereof, or a pharmaceutically acceptable salt of the compound or prodrug; or a cdk2, cdk5, and/or GSK-3 activity inhibiting amount of a pharmaceutical composition comprising a compound of formula (I), a prodrug thereof, or a pharmaceutically acceptable salt of the compound or prodrug, and a pharmaceutically acceptable carrier, vehicle, or diluent.

In another aspect, the invention provides pharmaceutical compositions comprising an amount of a compound of formula (I), a prodrug thereof, or a pharmaceutically acceptable salt of the compound or prodrug, optionally in combination with an amount of one or more of: (i) an anti-angiogenesis agent, (ii) a signal transduction inhibitor, (iii) an anti-proliferative agent, (iv) an NK-1 receptor antagonist, (v) a 5HT_{1D} receptor antagonist, (vi) a selective serotonin reuptake inhibitor (SSRI), (vii) an anti-psychotic agent, (viii) an acetylcholinesterase inhibitor, (ix) a neuroprotectant, (x) tissue plasminogen activator (TPA), (xi) neutrophil inhibitory factor (NIF), or a (xii) a potassium channel modulator.

In yet another aspect, the invention provides methods of treating cdk2, cdk5, and/or GSK-3 mediated conditions, diseases, or symptoms in a mammal in need of such treatment which methods comprise administering to the mammal: (i) a therapeutically effective amount of a compound of formula (I), a prodrug thereof, or a pharmaceutically acceptable salt of the compound or prodrug; (ii) a therapeutically effective amount of a pharmaceutical composition comprising a compound of formula (I), a prodrug thereof, or a pharmaceutically acceptable salt of the compound or prodrug, and a pharmaceutically acceptable carrier, vehicle, or diluent; (iii) a therapeutically effective amount of a combination of a compound of formula (I), a prodrug thereof, or a pharmaceutically acceptable salt of the compound or prodrug, and one or more of: (i) an anti-angiogenesis agent, (ii) a signal transduction inhibitor, (iii) an anti-proliferative agent, (iv) an NK-1 receptor antagonist, (v) a 5HT_{1D} receptor antagonist, (vi) a selective serotonin reuptake inhibitor (SSRI), (vii) an anti-psychotic agent, (viii) an acetylcholinesterase inhibitor, (ix) a neuroprotectant, (x) tissue plasminogen activator (TPA), (xi) neutrophil inhibitory factor (NIF), and (xii) a potassium channel modulator; or (iv) a therapeutically effective amount of a pharmaceutical composition comprising the aforementioned combinations.

Preferred conditions, diseases, and symptoms treatable according to the methods of the instant invention are those selected from the group consisting of Alzheimer's Disease, asthma, atherosclerosis, anxiety, bipolar disorder, cancer, diabetes, dementia, depression, frailty, hair loss, heart failure, essential hypertension, hyperglycemia, hyperlipidemia, hypoglycemia, inflammation, ischemia, male fertility and sperm motility, mood disorders, neuronal cell death, obesity, obsessive compulsive disorder, polycystic ovary disorder, schizophrenia, stroke, Syndrome X, and traumatic brain injury.

Frailty is characterized by the progressive loss of skeletal muscle mass resulting in a high risk of injury from fall, difficulty in recovery from illness, prolongation of hospitalization, and long-term disability requiring assistance in daily living. The reduction of muscle mass and physical strength typically leads to

5 diminished quality of life, loss of independence, and mortality. Frailty is normally associated with aging, but may also result when muscle loss and reduced strength occur due to other factors, such as disease-induced cachexia, immobilization, or drug-induced sarcopenia. Another term that has been used to denote frailty is sarcopenia, which is a generic term for the loss of skeletal muscle mass, or quality.

10 Examples of skeletal muscle properties that contribute to its overall quality include contractility, fiber size and type, fatiguability, hormone responsiveness, glucose uptake/metabolism, and capillary density.

Generally preferred anti-angiogenesis agents may comprise, for example, matrix metalloproteinase-2 (MMP-2) inhibitors, matrix metalloproteinase-9 (MMP-9)

15 inhibitors, and cyclooxygenase-II (COX-II) inhibitors. Examples of useful MMP-2 and MMP-9 inhibitors are disclosed in, for example, PCT International Application Publication Nos. WO 98/34915 and WO 98/34918, and U.S. Pat. Nos. 5,240,958; 5,310,763; 5,455,258; 5,506,242; 5,530,161; 5,552,419; 5,672,615; 5,861,510; 5,863,949; 5,932,595; 5,994,351; 6,077,864; 6,087,392; 6,090,852; 6,110,964;

20 6,147,061; 6,147,074; 6,303,636; 6,380,219; and 6,387,931. Examples of COX-II inhibitors useful in the present combinations and methods comprise CELEBREX® (celecoxib, U.S. Pat. No. 5,466,823), valdecoxib (U.S. Pat. No. 5,633,272), and rofecoxib (U.S. Pat. No. 5,474,995). Generally preferred MMP-2 and MMP-9 inhibitors are those exhibiting little or no activity inhibiting MMP-1. Especially preferred

25 MMP-2 and MMP-9 inhibitors are those that selectively inhibit MMP-2 and/or MMP-9 relative to other MMP inhibitors, i.e., MMP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13. Specific examples of MMP inhibitors useful in the present combinations and methods comprise AG-3340, RO 32-3555, RS 13-0830, and the following compounds:

30 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclopentyl)-amino]-propionic acid;

3-exo-3-[4-(4-fluoro-phenoxy)-benzenesulfonyl-amino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide;

- (2R, 3R) 1-[4-(2-chloro-4-fluoro-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide;
- 4-[4-(4-fluoro-phenoxy)-benzenesulfonyl-amino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
- 5 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-cyclobutyl)-amino]-propionic acid;
- 4-[4-(4-chloro-phenoxy)-benzenesulfonyl-amino]-tetrahydro-pyran-4-carboxylic acid hydroxyamide;
- (R)-3-[4-(4-chloro-phenoxy)-benzenesulfonyl-amino]-tetrahydro-pyran-3-
- 10 carboxylic acid hydroxyamide;
- (2R, 3R) 1-[4-(4-fluoro-2-methyl-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide;
- 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(1-hydroxycarbamoyl-1-methyl-ethyl)-amino]-propionic acid;
- 15 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(4-hydroxycarbamoyl-tetrahydro-pyran-4-yl)-amino]-propionic acid;
- 3-exo-3-[4-(4-chloro-phenoxy)-benzenesulfonyl-amino]-8-oxa-bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide;
- 3-endo-3-[4-(4-fluoro-phenoxy)-benzenesulfonyl-amino]-8-oxa-
- 20 bicyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; and
- (R)-3-[4-(4-fluoro-phenoxy)-benzenesulfonyl-amino]-tetrahydro-furan-3-carboxylic acid hydroxyamide; and the pharmaceutically acceptable salts and solvates thereof.

Generally preferred signal transduction inhibitors may comprise, for example,

25 epidermal growth factor receptor (EGFR) response inhibitors, such as EGFR antibodies, EGF antibodies, and molecules that are EGFR inhibitors; vascular endothelial growth factor (VEGF) inhibitors; and erbB2 receptor inhibitors, such as molecules or antibodies that bind to the erbB2 receptor, for example, HERCEPTIN® (Genentech Inc.; South San Francisco, CA). EGFR inhibitors are described in, for

30 example, PCT International Application Publication No. WO 98/14451, and U.S. Pat. Nos. 5,679,683; 5,747,498; and 6,391,874. EGFR-inhibiting agents may comprise, for example, the monoclonal antibodies C225 and anti-EGFR 22Mab (Imclone Systems, Inc.), ZD-1839, BIBX-1382, MDX-103, VRCTC-310, and EGF fusion toxin (Seragen Inc.; Hopkinton, MA). VEGF inhibitors are disclosed in, for example, PCT

International Application Publication No. WO 99/24440, and US. Pat. Nos. 5,792,783; 5,834,504; 5,851,999; 5,883,113; 5,886,020; 6,051,593; 6,114,371; 6,133,305; 6,162,804; 6,174,889; 6,207,669; 6,235,741; 6,291,455; 6,294,532; 6,310,238; 6,380,203; and 6,395,734. Specific VEGF inhibitors may comprise, for example, Su-
5 5416, IM862, anti-VEGF monoclonal antibody (Cytran Inc.; Kirkland, WA), and angiozyme (Ribozyme; Boulder, CO). ErbB2 receptor inhibitors are disclosed in, for example, PCT International Application Publication Nos. WO 97/13760, WO 99/35132, and WO 99/35146, and US. Pat. Nos. 5,679,683; 5,587,458; 5,877,305; 6,207,669; and 6,391,874. Specific erbB2 receptor inhibitors may comprise, for
10 example, GW-282974 (Glaxo Wellcome plc.), and the monoclonal antibody AR-209 (Aronex Pharmaceuticals Inc.; The Woodlands, TX).

Generally preferred anti-proliferative agents may comprise, for example, cytotoxic lymphocyte antigen 4 (CTLA4) antibodies, and other agents capable of blocking CTLA4; and farnesyl transferase inhibitors.

15 Examples of NK-1 receptor antagonists are disclosed in, for example, US. Pat. Nos. 5,122,525; 5,162,339; 5,232,929; 5,332,817; 5,703,240; 5,716,965; 5,719,147; 5,744,480; 5,763,699; 5,773,450; 5,807,867; 5,843,966; 5,852,038; 5,886,009; and 5,939,433.

Examples of 5HT_{1D} receptor antagonists useful in the present combinations
20 and methods are disclosed in, for example, PCT International Application Publication No. WO 94/21619, and U.S. Pat. Nos. 5,358,948; 5,510,350; 6,380,186; 6,403,592; 6,423,708; and 6,462,048.

Examples of SSRI's useful in the present combinations and methods may comprise, for example, fluoxetine (U.S. Pat. No. 4,314,081), paroxetine (U.S. Pat. No.
25 4,007,196), sertraline (U.S. Pat. No. 4,536,518), fluvoxamine (U.S. Pat. No. 4,085,225), venlafaxine hydrochloride (EFFEXOR[®], U.S. Pat. No. 4,535,186), nefazodone hydrochloride (SERZONE[®], U.S. Pat. No. 4,338,317), and bupropion hydrochloride (WELLBUTRIN[®], U.S. Pat. Nos. 3,819,706 and 3,885,046).

Generally preferred anti-psychotic agents useful in the present combinations
30 and methods may comprise, for example, ziprasidone (GEODON[®], U.S. Pat. No. 5,312,925), olanzapine (U.S. Pat. No. 5,229,382), risperidone (U.S. Pat. No. 4,804,663), L-745,870, sonopiprazole, RP-62203 (fananserine), NGD-941, balaperidone, flesinoxan (U.S. Pat. No. 4,833,142), and gepirone (U.S. Pat. No. 4,423,049).

Generally preferred acetylcholinesterase inhibitors useful in the present combinations and methods may comprise, for example, donepezil (ARICEPT[®], U.S. Pat. No. 4,895,841), rivastigmine (EXELON[®], U.S. Pat. No. 4,948,807), metrifonate (U.S. Pat. No. 2,701,225), galanthamine, physostigmine, tacrine, huperzine, and
5 icopezil (U.S. Pat. No. 5,538,984).

Generally preferred neuroprotectants useful in the instant combinations and methods may comprise, for example, NMDA receptor antagonists. Specific NMDA receptor antagonists comprise, for example, (1S, 2S)-1-(4-hydroxyphenyl)-2-(4-hydroxy-4-phenylpiperidin-1-yl)-1-propanol (U.S. Pat. No. 5,272,160); eliprodil (US.
10 Pat. No. 4,690,931); and gavestenel (U.S. Pat. No. 5,373,018). Examples of additional NMDA antagonists are disclosed in, for example, U.S. Pat. Nos. 4,690,931; 5,185,343; 5,272,160; 5,356,905; 5,373,018; 5,744,483; 5,962,472; 6,046,213; 6,124,317; 6,124,323; 6,130,234; 6,218,404; 6,333,036; and 6,448,270; and in PCT International Application Publication Nos. WO 97/23202 and WO 98/18793.

15 A generally preferred potassium channel modulator comprises, for example, BMS-204352 (flindokaliner, U.S. Pat. No. 5,602,169).

The disclosures of all of the above U.S. patents are incorporated herein in their entirety by reference.

The compounds of formula (I), the prodrugs thereof, and the pharmaceutically
20 acceptable salts of the compounds and prodrugs, may be administered to a mammal at dosage levels in the range of from about 0.0001 mg to about 1,000 mg per day. For a normal adult human having a body mass of about 70 kg, a dosage in the range of from about 0.01 mg to about 500 mg per kg body mass is typically sufficient. However, some variability in the general dosage range may be required depending
25 upon the age and mass of the subject being treated, the intended route of administration, the particular compound being administered, and the like. The determination of dosage ranges and optimal dosages for a particular mammalian subject is within the ability of one of ordinary skill in the art having benefit of the instant disclosure.

30 According to the methods of the present invention, the compounds of formula (I), the prodrugs thereof, and the pharmaceutically acceptable salts of the compounds and prodrugs, or the aforementioned combinations thereof, are preferably administered in the form of a pharmaceutical composition comprising a pharmaceutically acceptable carrier, vehicle, or diluent. Accordingly, an amount of a

compound of formula (I), a prodrug thereof, or a pharmaceutically acceptable salt of the compound or prodrug, or the aforementioned combinations, may be administered to a subject separately, or together, in any conventional oral, rectal, transdermal, parenteral (e.g., intravenous, intramuscular, or subcutaneous), intracisternal, 5 intravaginal, intraperitoneal, intravesical, local (e.g., powder, ointment, or drop), or buccal, or nasal dosage form.

Pharmaceutical compositions suitable for parenteral injection may comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions, or emulsions, and sterile powders for extemporaneous reconstitution 10 into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, vehicles, and diluents include water, ethanol, polyols (such as propylene glycol, polyethylene glycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as 15 lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

The pharmaceutical compositions of the invention may further comprise adjuvants, such as preserving, wetting, emulsifying, and dispersing agents. Prevention of microorganism contamination of the instant compositions can be 20 accomplished with various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example, sugars, sodium chloride, and the like. Prolonged absorption of injectable pharmaceutical compositions may be effected by the use of agents capable of delaying absorption, for example, aluminum monostearate and 25 gelatin.

Solid dosage forms for oral administration include capsules, tablets, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert conventional pharmaceutical excipient (or carrier) such as sodium citrate or dicalcium phosphate, or (a) fillers or extenders, as for example, starches, 30 lactose, sucrose, mannitol, and silicic acid; (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia; (c) humectants, as for example, glycerol; (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid certain complex silicates, and sodium carbonate; (e) solution retarders, as for example, paraffin; (f)

absorption accelerators, as for example, quaternary ammonium compounds; (g) wetting agents, as for example, cetyl alcohol and glycerol monostearate; (h) adsorbents, as for example, kaolin and bentonite; and/or (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules and tablets, the dosage forms may further comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft or hard filled gelatin capsules using such excipients as lactose or milk sugar, as well as high molecular weight polyethylene glycols, and the like.

Solid dosage forms such as tablets, dragees, capsules, and granules can be prepared with coatings and shells, such as enteric coatings and others well-known to one of ordinary skill in the art. They may also comprise opacifying agents, and can also be of such composition that they release the active compound(s) in a delayed, sustained, or controlled manner. Examples of embedding compositions that can be employed are polymeric substances and waxes. The active compound(s) can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage form may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil, and sesame seed oil, glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, or mixtures of these substances, and the like.

Besides such inert diluents, the pharmaceutical composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.


Suspensions, in addition to the active compound(s), may further comprise suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, or mixtures of the aforementioned substances, and the like.

Compositions for rectal or vaginal administration preferably comprise suppositories, which can be prepared by mixing an active compound(s) with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax, which are solid at ordinary room temperature, but liquid at body temperature, and therefore, melt in the rectum or vaginal cavity thereby releasing the active component.

Dosage forms for topical administration may comprise ointments, powders, sprays and inhalants. The active agent(s) are admixed under sterile condition with a pharmaceutically acceptable carrier, vehicle, or diluent, and any preservatives, buffers, or propellants that may be required.

The compounds of formula (I), the prodrugs thereof, and the pharmaceutically acceptable salts of the compounds and prodrugs, may be prepared according to the exemplary synthetic routes disclosed in the Schemes and Examples hereinbelow, as well as by other conventional organic preparative methods known, or apparent in light of the instant disclosure, to one of ordinary skill in the relevant art. It is to be understood that the methods disclosed in the instant Schemes are intended for purposes of exemplifying the instant invention, and are not to be construed in any manner as limitations thereon.


Scheme 1



In Scheme 1, an appropriately-substituted cyclobutanone derivative (1), prepared as disclosed in the aforementioned PCT International Application Publication No. WO 02/18346, is treated with a reducing agent, preferably sodium

borohydride, in a suitable solvent, such as a mixture of tetrahydrofuran (THF) and ethanol (EtOH), at below ambient temperature, preferably about -78°C , followed by warming to 0°C , or ambient temperature. It has been found that the *cis* isomer of the resulting cyclobutanol (2) is formed predominantly over the corresponding *trans* isomer, typically in ratios of $>10:1$. Cyclobutanol (2) is treated with a carbonic acid equivalent, preferably triphosgene or 1,1'-carbonyldiimidazole (CDI), to form the activated intermediate (3), in a solvent such as ethyl acetate (EtOAc) or methylene chloride, at or below ambient temperature. An amine base, preferably pyridine, is added to reactions utilizing triphosgene, and may optionally be employed for reactions using CDI. Addition of an appropriately-substituted amine HNR^2R^3 to the solution of (3), typically at a temperature of between ambient temperature and the reflux temperature of the solvent employed, affords protected pyrazole (4). The *tert*-butyl protecting group is cleaved by treating (4) with trifluoroacetic acid (TFA), at elevated temperature, preferably $>70^{\circ}\text{C}$, to afford (I).


Alternatively, the compounds of formula (I) may be prepared according to the method disclosed in Scheme 2.



In Scheme 2, the *tert*-butyl protecting group of compound (2) is cleaved by treatment with an acid, preferably TFA, at elevated temperature. The resulting ester (5) is cleaved by treatment with an aqueous base, such as ammonium hydroxide or

sodium hydroxide, to provide alcohol (6). Alcohol (6) is then treated with CDI in an organic solvent, preferably EtOAc, to afford intermediate imidazolidine (7) which, if desired, may be isolated by conventional workup. The reaction of an appropriately-substituted amine HNR^2R^3 with (7) is effected in an organic solvent, preferably EtOAc, provides (I). Where needed, an amine base, such as triethylamine (Et_3N), 4-(dimethylamino)pyridine (DMAP), or a polymer-supported DMAP derivative, may be added. When an elevated temperature is required, such temperatures may be achieved by known methods, including heating the reaction in a microwave apparatus.

Alternatively, the compounds of formula (I) may be prepared according to the method disclosed in Scheme 3.




In Scheme 3, the *tert*-butyl protecting group of compound (1) is removed by treatment with acid as described hereinabove in Scheme 2. Deprotected pyrazole (8) is then treated with di-*tert*-butyldicarbonate in the presence of a base, preferably Et_3N and DMAP, in an aprotic solvent such as methylene chloride, at ambient temperature. The resulting Boc-protected pyrazole (9) is isolated as a mixture of carbamate isomers that may be employed subsequently without further purification. Pyrazole (9) is treated with a reducing agent, preferably sodium borohydride, in a suitable solvent, such as a mixture of THF and EtOH, at or below ambient temperature, to afford alcohol (10). Reaction of (10) with an appropriately-substituted amine HNR^2R^3 affords (11) which is then deprotected by treatment with an acid, such as TFA or,

alternatively, by warming a solution of (11) in acetonitrile or dimethylsulfoxide (DMSO) in a microwave apparatus at about 150° C.

Alternatively, the compounds of formula (I) may be according prepared according to the method disclosed in Scheme 4.

5

Scheme 4



In Scheme 4, compound (12), prepared as disclosed in the aforementioned PCT International Application Publication No. WO 02/18346, is treated with an excess of di-*tert*-butyldicarbonate in the presence of a base, preferably Et_3N and DMAP, in an aprotic solvent such as methylene chloride, at ambient temperature, to provide the *bis*-carbamoylated product (13). Treatment of (13) with aqueous acid, preferably *p*-toluenesulfonic acid, in a mixture of water and acetone, preferably at reflux temperature, affords ketone (14). Ketone (14) is then treated with a reducing agent, preferably sodium borohydride, in a suitable solvent, such as a mixture of THF and EtOH, at below ambient temperature, preferably -78° C, followed by warming to

0° C, or ambient temperature. As in Scheme 1, it has been found that the *cis* isomer of the resulting cyclobutanol (15) is formed predominantly over the corresponding *trans* isomer, typically in ratios of ~10:1. Cyclobutanol (15) is then treated with a carbonic acid equivalent, preferably triphosgene or CDI, in a solvent such as EtOAc or methylene chloride, at or below ambient temperature. An amine base, preferably pyridine, is added to reactions utilizing triphosgene, and may optionally be employed for reactions using CDI. Addition of an appropriately-substituted amine HNR^2R^3 , typically at a temperature of between ambient temperature and the reflux temperature of the solvent employed, affords protected pyrazole (16). The *tert*-butyl carbamate protecting groups are cleaved by treating (16) with trifluoroacetic acid (TFA) at ambient temperature to afford amine (17). Amine (17) is then treated under standard acylation conditions, with either a carboxylic acid and an amine coupling reagent, or a carboxylic acid chloride, and a base, such as Et_3N or pyridine, to afford (4). The *tert*-butyl protecting group is cleaved by treating (4) with trifluoroacetic acid (TFA), at elevated temperature, preferably >70° C, to furnish compound (I).

PREPARATIVE EXPERIMENTAL

Unless otherwise noted, all reagents employed were obtained commercially. Unless otherwise noted, the following experimental abbreviations have the meanings indicated:

DMF - dimethylformamide
Et₃SiH - triethylsilane
HPLC - high performance liquid chromatography
h - hour(s)
M - molar
MeOH - methanol
min - minute(s)
IPA - isopropanol
mL - milliliter(s)
mmol - millimole(s)
HPLC – high performance liquid chromatography
MS - mass spectrometry

Unless otherwise noted, preparation of the various N-(5-cyclobutyl-1H-pyrazol-3-yl)-amide starting materials from 2-*tert*-butyl-5-(3,3-dimethoxy-cyclobutyl)-

2H-pyrazol-3-ylamine was carried out according to the methods described in the
aforementioned PCT International Application Publication No. WO02/18346. 2-
Methyl-tetrahydro-furan-2-carboxylic acid and 4-methyl-tetrahydro-pyran-4-carboxylic
acid were prepared from tetrahydro-furan-2-carboxylic acid methyl ester and
5 tetrahydro-pyran-4-carboxylic acid methyl ester, respectively, according to the
method of Regan (J. Med. Chem., 45, 2994-3008 (2002)). Carboxylic acid chlorides
that were not commercially available were prepared from the corresponding
carboxylic acids by treatment with thionyl chloride (Org. Syn., Coll. Vol., 3, 169
(1955)).

10 Preparative reversed-phase HPLC purifications were carried out on a system
obtained from Shimadzu Scientific Instruments, Inc.; Columbia, MD (Model LC-8A
Prep LC, SPD-10A UV-vis detector, FRC-10A fraction collector, SIL-10AP auto-
injector). All microwave chemistry was performed using an Emrys Optimizer®
(Personal Chemistry Inc.; Foxboro, MA).

15 Preparation 1

2-Methyl-2-pyridin-2-yl-propionic acid ethyl ester and 2-Pyridin-2-yl-propionic acid
ethyl ester

A solution of *n*-butyllithium in hexanes (2.5 M, 121 mL) was added slowly to a
solution of diisopropylamine (42 mL) in THF (120 mL) at -78 °C, and the resulting
20 solution was stirred for 15 min. Pyridin-2-yl-acetic acid ethyl ester (9.2 mL) was then
added and the mixture was stirred for 30 min before iodomethane (18.9 mL) was
added. The reaction mixture was stirred at -78 °C for 15 min and then at room
temperature for 3 h. The reaction mixture was diluted with water and extracted with
CH₂Cl₂ (3x). After concentration, the residue was purified by silica gel
25 chromatography to afford the title compounds separately. 2-Methyl-2-pyridin-2-yl-
propionic acid ethyl ester: MS (M+H)⁺ = 194.1. 2-Pyridin-2-yl-propionic acid ethyl
ester: MS (M+H)⁺ = 180.1.

Example 1

Benzyl-carbamic acid *cis*-3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-
30 yl]-cyclobutyl ester

Step A

A solution of trimethylaluminum in heptane (2.0 M, 19.5 mL) was added slowly
to a solution of 2-*tert*-butyl-5-(3,3-dimethoxy-cyclobutyl)-2H-pyrazol-3-ylamine (8.98
g) in CH₂Cl₂ (60 mL) at room temperature. After 15 min, a solution of 2-methyl-2-

pyridin-2-yl-propionic acid ethyl ester (6.85 g) in CH_2Cl_2 (60 mL) was added. The reaction mixture was heated to reflux overnight. Saturated aqueous NH_4Cl solution was added and the mixture was extracted with CH_2Cl_2 (2x). The combined organic layers were dried over Na_2SO_4 and concentrated to afford N-[2-tert-butyl-5-(3,3-dimethoxy-cyclobutyl)-2H-pyrazol-3-yl]-2-pyridin-2-yl-isobutyramide as an oil that was used without further purification.

Step B

A mixture of the product of Step A (1.0 g) and *p*-toluenesulfonic acid monohydrate (50 mg) in water (0.7 mL) and acetone (9 mL) was heated at reflux for 7 h. The solution was cooled, concentrated, and the resulting residue was partitioned between EtOAc and saturated aqueous NaHCO_3 solution. The organic layer was dried over Na_2SO_4 and concentrated to afford the product ketone that was used without further purification.

Step C

Sodium borohydride (118 mg) was added to a solution of the product of Step B (1 g) in THF (9 mL) and EtOH (1.3 mL) at -78°C . The reaction mixture was stirred for 15 min at -78°C and then at 0°C , and saturated aqueous NH_4Cl solution was added to quench excess hydride. The mixture was extracted with CH_2Cl_2 (3x) and the combined organic layers were dried over Na_2SO_4 and concentrated. The alcohol product was purified by silica gel chromatography.

Step D

To a solution of the product of Step C (50 mg) in CH_2Cl_2 (0.5 mL) at 0°C was added sequentially a solution of triphosgene (29 mg) in CH_2Cl_2 (0.5 mL) and pyridine (0.036 mL). The mixture was stirred at room temperature for 1 h, and benzylamine (45 mg) was added. After 3 h, saturated aqueous NH_4Cl solution was added and the mixture was extracted with CH_2Cl_2 (3x). The combined organic layers were dried over Na_2SO_4 and concentrated. The residue was purified by silica gel chromatography to afford the product carbamate.

Step E

A solution of the product of Step E (48 mg), Et_3SiH (0.046 mL), and TFA (1 mL) was heated to reflux for 12 h. The solution was concentrated and the residue partitioned between saturated aqueous NaHCO_3 solution and CH_2Cl_2 . The organic layer was dried over Na_2SO_4 and concentrated. The residue was purified by silica gel chromatography to afford the title product as a white solid. MS $(\text{M}+\text{H})^+ = 434.2$.

The following compounds were prepared in a manner analogous to that described in Example 1 using appropriate starting materials.

Example	Name	MS (M+H) ⁺
2	Benzyl-methyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	448.2
3	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	400.2

The following compounds were prepared in a manner analogous to that described in Example 1, Steps C to E, using appropriate starting materials.

Example	Name	MS (M+H) ⁺
4	Benzyl-ethyl-carbamic acid <i>cis</i> -3-{5-[[<i>(R)</i> -(tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	413.4
5	Methyl-pyridin-3-ylmethyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-1H-pyrazol-3-yl)-cyclobutyl ester	372.4
6	1,3-Dihydro-isoindole-2-carboxylic acid <i>cis</i> -3-(5-isobutyrylamino-1H-pyrazol-3-yl)-cyclobutyl ester	369.4
7	Methyl-phenyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-1H-pyrazol-3-yl)-cyclobutyl ester	357.4
8	Phenyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-1H-pyrazol-3-yl)-cyclobutyl ester	343.4

Example 9

Cyclohexylmethyl-carbamic acid *cis*-3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester

10 Step A

A solution of *N*-[2-*tert*-butyl-5-(*cis*-3-hydroxy-cyclobutyl)-2H-pyrazol-3-yl]-3-pyrazol-1-yl-propionamide (200 mg, prepared as in Example 1, Step C using appropriate starting materials) and CDI (100 mg) in EtOAc (3 mL) was stirred at room temperature. After 45 min, cyclohexyl-methylamine (0.12 mL) was added and the solution was heated at reflux overnight. The solution was cooled and then diluted with EtOAc and washed sequentially with portions of saturated aqueous NH₄Cl solution,

saturated aqueous NaHCO₃ solution, and saturated aqueous NaCl solution. The organic layer was dried over Na₂SO₄ and then concentrated. The residue was purified by silica gel chromatography to afford the product carbamate as a white solid.

Step B

- 5 A solution of cyclohexylmethyl-carbamic acid *cis*-3-[1-*tert*-butyl-5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester from Step A (122 mg) in TFA (2 mL) was heated at 120 °C in a microwave apparatus for 10 min. The solution was concentrated and the residue was dissolved in EtOAc. The resulting solution was washed sequentially with portions of saturated aqueous NaHCO₃ solution and
- 10 saturated aqueous NaCl solution. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by silica gel chromatography to afford the title product as a white solid. MS (M+H)⁺ = 415.5.

- The following compounds were prepared in a manner analogous to that described in Example 9 using appropriate starting materials. For Examples
- 15 designated with an asterisk, Step B was conducted in TFA at reflux overnight.

Example	Name	MS (M+H) ⁺
10	Diethyl-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	363.4
11	(2-Methoxy-1-methyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	379.4
12	(2-Methoxy-ethyl)-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	365.4
13	(2-Dimethylamino-ethyl)-methyl-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	392.4
14	Isobutyl-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	363.4
15	Benzyl-methyl-carbamic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	413.4
16	(2-Methoxy-ethyl)-methyl-carbamic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	381.4
	Benzyl-carbamic acid <i>cis</i> -3-[5-	

17*	[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	399.4
18*	Piperidine-1-carboxylic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	377.4
19	Isopropyl-methyl-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	363.4
20	(2-Pyridin-3-yl-ethyl)-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	412.3
21*	(2-Chloro-benzyl)-carbamic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	433.3
22*	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	400.4
23*	(3-Chloro-benzyl)-carbamic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	433.3
24	Phenethyl-carbamic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	413.4
25	Pyridin-3-ylmethyl-carbamic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	400.4
26	3,4-Dihydro-1H-isoquinoline-2-carboxylic acid <i>cis</i> -3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	425.6
27	Benzyl-methyl-carbamic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	423.5
28	Benzyl-carbamic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	409.4
29	Methyl-propyl-carbamic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	375.4
30*	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	410.4

31	Propyl-carbamic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	361.4
32	Benzyl-carbamic acid <i>cis</i> -3-[5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	385.4
33	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	386.4
34	Isobutyl-carbamic acid <i>cis</i> -3-[5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	351.4
35	Benzyl-methyl-carbamic acid <i>cis</i> -3-[5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	399.4
36*	[2-(3,5-Dimethyl-pyrazol-1-yl)-ethyl]-carbamic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	441.4
37*	Benzyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-1H-pyrazol-3-yl)-cyclobutyl ester	357.4
38	(Tetrahydro-furan-2-ylmethyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-1H-pyrazol-3-yl)-cyclobutyl ester	351.4
39	(2-Phenoxy-ethyl)-carbamic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	439.4
40	3,4-Dihydro-1H-isoquinoline-2-carboxylic acid <i>cis</i> -3-[5-(3-pyrazol-1-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	435.4
41*	Dimethyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	372.3

Example 42

Ethyl-carbamic acid *cis*-3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester

5 Step A

Sodium borohydride (330 mg) was added to a solution of *N*-[2-*tert*-butyl-5-(3-oxo-cyclobutyl)-2H-pyrazol-3-yl]-2-phenyl-isobutyramide (2.8 g) in THF (100 mL) and EtOH (10 mL) at -60 °C. The mixture was warmed to room temperature, and held at

that temperature for 1.5 h. Excess hydride was quenched by the addition of MeOH (50 mL). The mixture was concentrated and the resulting residue was dissolved in EtOAc. The solution was washed sequentially with saturated aqueous K_2CO_3 solution (2x) and saturated aqueous NaCl solution. The organic layer was dried over Na_2SO_4 and concentrated to afford the product alcohol as a light yellow solid that was used without further purification.

Step B

A solution of the product of Step A (2.8 g) in TFA (50 mL) was heated to reflux for 48 h. The solution was concentrated and the resulting residue was dissolved in EtOAc. The solution was washed sequentially with saturated aqueous $NaHCO_3$ solution and saturated aqueous NaCl solution. The organic layer was dried over Na_2SO_4 and concentrated to afford the product ester as a residue that was used without further purification.

Step C

A solution of the product of Step B (1.9 g), concentrated aqueous NH_4OH (10 mL), and MeOH (20 mL) was heated to 65 °C for 90 min. The solution was concentrated, the resulting residue was diluted with EtOAc, and the solution washed sequentially with portions of saturated aqueous $NaHCO_3$ solution and saturated aqueous NaCl solution. The organic layer was dried over Na_2SO_4 and concentrated to afford the product alcohol as a solid that was used without further purification.

Step D

A mixture of the product of Step C (1.1 g) and CDI (714 mg) in EtOAc (30 mL) was stirred for 3 h. The solution was diluted with EtOAc and washed sequentially with portions of saturated aqueous $NaHCO_3$ solution and saturated aqueous NaCl solution. The organic layer was dried over Na_2SO_4 and concentrated to afford the product imidazolidine as a solid that was used without further purification.

Step E

A mixture of the product of Step D (100 mg), PS-DMAP resin (0.15 mmol; Argonaut Technologies; Foster City, CA), and ethylamine hydrochloride (27 mg) in EtOAc (2 mL) was heated at 35 °C for 10 h. The resin was removed by filtration and the filtrate was concentrated. The resulting residue was purified by reversed-phase preparative HPLC to afford the title product as a white solid. MS $(M+H)^+ = 371.2$.

The following compounds were prepared in a manner analogous to that described in Example 42 using appropriate starting materials. For Examples 43 to 71, DMAP was used in Step E, rather than PS-DMAP resin.

Example	Name	MS (M+H) ⁺ or MS (M-H) ⁻
43	(2-Cyano-ethyl)-methyl-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	374.4 (+)
44	(3-Chloro-benzyl)-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester	431.4 (+)
45	(2-Chloro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	419.4 (+)
46	(3-Chloro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	419.4 (+)
47	(4-Chloro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	419.4 (+)
48	(4-Methoxy-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	415.4 (+)
49	(2,2-Dimethyl-propyl)-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	365.5 (+)
50	(2-Phenylamino-ethyl)-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	414.4 (+)
51	Cyclohexylmethyl-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	391.5 (+)
52	(2-Isopropoxy-ethyl)-carbamic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	381.4 (+)
53	3,4-Dihydro-1H-isoquinoline-2-carboxylic acid <i>cis</i> -3-{5-[(<i>(R)</i>)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl}-cyclobutyl ester	411.3 (+)
	(4-Isopropyl-benzyl)-carbamic acid <i>cis</i> -	

54	3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl)-cyclobutyl ester	427.4 (+)
55	(2-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl)-cyclobutyl ester	403.3 (+)
56	Cyclopropylmethyl-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl)-cyclobutyl ester	349.3 (+)
57	(2-Phenyl-propyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl)-cyclobutyl ester	413.4 (+)
58	(4-Cyano-cyclohexylmethyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl)-cyclobutyl ester	416.4 (+)
59	(2-Piperidin-1-yl-ethyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	406.3 (+)
60	(2-Chloro-6-fluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	437.2 (+)
61	(2,4-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	421.2 (+)
62	(2,3-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	421.2 (+)
63	(2-Methyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	399.2 (+)
64	(2,6-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	421.2 (+)
65	(2-Fluoro-3-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	471.2 (+)
66	<i>tert</i> -Butyl-carbamic acid <i>cis</i> -3-{5-[(R)-tetrahydro-furan-2-carbonyl]-amino}-1H-pyrazol-3-yl)-cyclobutyl ester	351.3 (+)

67	(2-Phenoxy-ethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	415.2 (+)
68	(2-Methoxy-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	415.2 (+)
69	(2-Trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	453.2 (+)
70	(2-Fluoro-5-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	471.2 (+)
71	(2,5-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	421.2 (+)
72	(1H-Benzoimidazol-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	423.1 (-)
73	(2-Pyridin-2-yl-ethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	398.2 (-)
74	(2-Pyrrolidin-1-yl-ethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	392.4 (+)
75	(1-Ethyl-pyrrolidin-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	406.4 (+)
76	(2-Morpholin-4-yl-ethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	408.3 (+)
77	[2-(<i>cis</i> -3-{5-[[((R)-Tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutoxycarbonylamino)-ethyl]-carbamic acid <i>tert</i> -butyl ester	438.4 (+)
78	(2-Dimethylamino-ethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	366.3 (+)
79	(2-Diethylamino-ethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-	394.3 (+)

	carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	
80	[2-(2-Methyl-5-nitro-imidazol-1-yl)-ethyl]-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	448.2 (+)
81	[2-(<i>cis</i> -3-{5-[(<i>R</i>)-Tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutoxycarbonylamino)-ethyl]-carbamic acid benzyl ester	472.2 (+)
82	(2-Acetylamino-ethyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	378.1 (-)
83	[2-(5-Nitro-pyridin-2-ylamino)-ethyl]-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	460.2 (+)
84	(2-Fluoro-6-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	471.2 (+)
85	(5-Methyl-furan-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	389.3 (+)
86	(2,2-Dimethyl-[1,3]dioxolan-4-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	409.3 (+)
87	Thiophen-2-ylmethyl-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	391.2 (+)
88	[2-(Pyridin-2-ylamino)-ethyl]-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	413.2 (-)
89	[2-(4-Trifluoromethyl-pyridin-2-ylamino)-ethyl]-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	483.2 (+)
90	(1-Methyl-1H-imidazol-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	389.4 (+)
91	(7-Methyl-imidazo[1,2- <i>a</i>]pyridin-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl)-amino]-	437.2 (-)

	2H-pyrazol-3-yl)-cyclobutyl ester	
92	(2-Imidazol-1-yl-ethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	389.4 (+)
93	(1H-Indol-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	424.3 (+)
94	[2-(2-Oxo-pyrrolidin-1-yl)-ethyl]-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	406.3 (+)
95	(2-Chloro-6-fluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	451.1 (+)
96	(2-Methyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	413.2 (+)
97	(4-Chloro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	433.2 (+)
98	(2-Fluoro-3-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	485.2 (+)
99	(4-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	403.2 (+)
100	(3,4-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	421.2 (+)
101	(4-Fluoro-2-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	471.2 (+)
102	(2,4,5-Trifluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	439.2 (+)
103	(4-Fluoro-3-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[[((R)-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl)-cyclobutyl ester	471.2 (+)
	(3,4,5-Trifluoro-benzyl)-carbamic acid	

104	<i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	439.2 (+)
105	(2,5-Dichloro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	453.1 (+)
106	(3,4-Dichloro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	453.1 (+)
107	(5-Chloro-2-methyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	433.3 (+)
108	(2-Chloro-6-fluoro-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	409.1 (+)
109	(2-Trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	425.1 (+)
110	(2-Fluoro-6-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	443.2 (+)
111	(2,6-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	393.2 (+)
112	(2-Chloro-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	391.2 (+)
113	(4-Chloro-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	391.2 (+)
114	(2-Phenyl-propyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	385.3 (+)
115	(2-Fluoro-3-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	443.3 (+)
116	(2,4-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	393.1 (+)
117	(2-Phenoxy-ethyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	387.1 (+)
118	(2-Phenylamino-ethyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	386.3 (+)

119	Benzyl-methyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	371.2 (+)
120	Cyclohexylmethyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	363.3 (+)
121	[2-(2-Methyl-pyridin-3-yl)-ethyl]-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	414.3 (+)
122	[2-(6-Methyl-pyridin-3-yl)-ethyl]-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	414.3 (+)
123	(3-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	403.2 (+)
124	(3,5-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	421.2 (+)
125	(3-Fluoro-5-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	471.1 (+)
126	[2-(6-Methoxy-pyridin-3-yl)-ethyl]-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	430.2 (+)
127	(3-Chloro-4-fluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(<i>R</i>)-tetrahydro-furan-2-carbonyl]-amino}-2H-pyrazol-3-yl}-cyclobutyl ester	437.4 (+)
128	Dimethyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	293.2 (-)
129	Methyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	281.3 (+)
130	Isopropyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	309.4 (+)
131	Piperidine-1-carboxylic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	335.4 (+)
132	Pyrrolidine-1-carboxylic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	321.4 (+)
	Butyl-carbamic acid <i>cis</i> -3-(5-	

133	isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	323.3 (+)
134	Propyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	309.2 (+)
135	(2,2-Dimethyl-propyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	337.3 (+)
136	Isobutyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	323.3 (+)
137	Morpholine-4-carboxylic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	337.2 (+)
138	(Tetrahydro-pyran-4-ylmethyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	365.4 (+)
139	(2,2-Dimethyl-tetrahydro-pyran-4-yl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	379.4 (+)
140	(Tetrahydro-furan-3-yl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	337.3 (+)
141	Cyclohexyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	349.3 (+)
142	(2-Methyl-butyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	337.3 (+)
143	Cyclopentyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	335.3 (+)
144	(2-Ethyl-butyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	351.3 (+)
145	Cyclobutyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	321.2 (+)
146	Azetidine-1-carboxylic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	307.2 (+)
147	(3,4,5,6-Tetrahydro-2H-[1,2']bipyridinyl-3-ylmethyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	441.2 (+)
148	(2,2-Diphenyl-ethyl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	447.2 (+)

149	(1-Benzyl-piperidin-4-yl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	440.2 (+)
150	((S)-1-Benzyl-pyrrolidin-3-yl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	426.2 (+)
151	((R)-1-Benzyl-pyrrolidin-3-yl)-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	426.2 (+)
152	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	434.1 (+)
153	Morpholine-4-carboxylic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	413.2 (+)
154	Dimethyl-carbamic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	371.2 (+)
155	Methyl-carbamic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	357.2 (+)
156	Ethyl-methyl-carbamic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	385.2 (+)
157	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	399.2 (+)
158	Ethyl-carbamic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	371.2 (+)
159	Propyl-carbamic acid <i>cis</i> -3-[5-(2-phenyl-butyrylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	385.2 (+)
160	Propyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	385.2 (+)
161	(Tetrahydro-pyran-4-ylmethyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	441.4 (+)
162	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	434.3 (+)
163	Ethyl-methyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	385.4 (+)
164	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	399.2 (+)

165	Benzyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	433.2 (+)
166	(2-Methoxy-benzyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	463.3 (+)
167	(2-Methyl-benzyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	447.4 (+)
168	(2-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	451.4 (+)
169	(2,6-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	469.4 (+)
170	(2-Trifluoromethyl-benzyl)-carbamic acid 3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	501.5 (+)
171	Ethyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-2H-pyrazol-3-yl)-cyclobutyl ester	295.3 (+)

Example 172

(Tetrahydro-pyran-4-ylmethyl)-carbamic acid *cis*-3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester

- 5 A solution of imidazole-1-carboxylic acid *cis*-3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester (200 mg) and C-(tetrahydro-pyran-4-yl)-methylamine (83 mg) in EtOAc was stirred at 70 °C overnight. After cooling, the title product was isolated by silica gel chromatography. MS (M+H)⁺ = 379.3.

- 10 The following compounds were prepared in a manner analogous to that described in Example 172 using appropriate starting materials. Amine starting materials that were obtained in the form of an acid addition salt were neutralized *in situ* by the addition of excess Et₃N. For Examples denoted with an asterisk, the reaction was heated in a microwave apparatus (150 °C, 35 min), rather than at reflux.

Example	Name	MS (M+H) ⁺ or MS (M-H) ⁻
173	Benzyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-	371.3 (+)

	3-yl]-cyclobutyl ester	
174*	(4-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	389.2 (+)
175	(4-Methoxy-benzyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	401.3 (+)
176*	(2-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	389.2 (+)
177*	(3-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	389.2 (+)
178	Benzyl-carbamic acid <i>cis</i> -3-{5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl}-cyclobutyl ester	409.2 (-)
179	(2-Chloro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl}-cyclobutyl ester	443.2 (-)
180	(2-Methoxy-2-methyl-propyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	367.3 (+)
181	(3,4-Dihydro-2H-pyran-2-ylmethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	377.3 (+)
182	(2-Methoxy-benzyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	401.3 (+)
183	(3-Methoxy-benzyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	401.3 (+)
184	(2-Methoxy-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	339.2 (+)
185	Butyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	337.3 (+)
186	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	337.3 (+)
187	Benzyl-methyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	385.3 (+)
	(Tetrahydro-furan-2-ylmethyl)-	

188	carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	365.2 (+)
189	(2-Phenylamino-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	400.3 (+)
190	(2,4-Dichloro-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	479.1 (+)
191	(2-Trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	479.2 (+)
192	(2,4-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	447.2 (+)
193	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	412.2 (+)
194	(R)-3-Propoxy-pyrrolidine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	393.3 (+)
195	Isochroman-1-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	427.3 (+)
196	[2-(2-Chloro-phenoxy)-propyl]-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	449.2 (+)
197	(2,2-Dimethyl-propyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	351.2 (+)
198	Piperazine-1,4-dicarboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester ethyl ester	422.2 (+)
199	4-Pyridin-2-yl-piperazine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	427.2 (+)
200	(2R,6S)-2,6-Dimethyl-morpholine-4-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-	379.2 (+)

	cyclobutyl ester	
201	(S)-3-Methoxy-pyrrolidine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	365.2 (+)
202	(2-Trifluoromethoxy-benzyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	455.1 (+)
203	4-Phenyl-piperazine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	426.3 (+)
204	3-Pyridin-2-yl-pyrrolidine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	412.3 (+)
205	[2-(Pyridin-3-yloxy)-propyl]-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	416.3 (+)
206	3-(Acetyl-methyl-amino)-pyrrolidine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	406.2 (+)
207	(4-Fluoro-2-trifluoromethyl-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	497.0 (+)
208	(3,4-Dichloro-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	480.2 (+)
209	(3,5-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	447.3 (+)
210	(3-Chloro-2-methyl-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	459.2 (+)
211	(4-Isopropyl-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	453.3 (+)
212	(2,6-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	447.2 (+)
	(3-Chloro-benzyl)-carbamic acid <i>cis</i> -	

213	3-{5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl}-cyclobutyl ester	445.1 (+)
214	(3-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-{5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl}-cyclobutyl ester	429.2 (+)
215	(5,7-Dimethyl-benzothiazol-2-yl)-carbamic acid <i>cis</i> -3-{5-[(1-methyl-cyclohexanecarbonyl)-amino]-1H-pyrazol-3-yl}-cyclobutyl ester	482.2 (+)
216	Indan-1-yl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	397.2 (+)
217	(1,2,3,4-Tetrahydro-naphthalen-1-yl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	411.2 (+)
218	(1-Phenyl-propyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	399.2 (+)
219	((S)-1-Phenyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	385.2 (+)
220	((R)-1-Phenyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	385.2 (+)

Example 221

Methyl-pyridin-3-ylmethyl-carbamic acid *cis*-3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester

5 Step A

A solution of *N*-[2-*tert*-butyl-5-(3-oxo-cyclobutyl)-2H-pyrazol-3-yl]-2,2-dimethyl-propionamide (6.9 g) in TFA (47 mL) was heated at reflux for 29 h. The solution was cooled, concentrated, and the resulting residue was partitioned between CH₂Cl₂ and saturated aqueous NaHCO₃ solution. The organic layer was dried over Na₂SO₄ and
10 concentrated to afford the product as a light yellow solid that was used without further purification.

Step B

A mixture of the product of Step A (2.7 g), di-*tert*-butyldicarbonate (3.6 g), Et₃N (2.85 mL), and DMAP (63 mg) in CH₂Cl₂ was stirred at room temperature for 2

h. The solution was concentrated and the resulting residue dissolved in EtOAc. The solution was washed sequentially with saturated aqueous NH_4Cl solution, water, and saturated aqueous NaCl solution. The organic layer was dried over Na_2SO_4 and concentrated to afford a mixture of carbamate isomers that was used without further purification.

Step C

Sodium borohydride (430 mg) was added to a solution of the product of Step B (4.8 g) in THF (34 mL) and EtOH (4.9 mL) at -78°C . After 10 min, the reaction mixture was warmed to 0°C for 15 min. Excess hydride was quenched by the addition of saturated aqueous NH_4Cl solution and the mixture was concentrated. The residue was dissolved in EtOAc and washed sequentially with saturated aqueous NaHCO_3 solution (2x), water, and saturated aqueous NaCl solution. The organic layer was dried over Na_2SO_4 and then was concentrated to afford the product as a yellow foam that was used without further purification.

Step D

The product of Step C (200 mg) was dissolved in CH_2Cl_2 and the resulting solution was cooled to 0°C . A solution of triphosgene (123 mg) in CH_2Cl_2 (0.5 mL) was added followed by pyridine (0.125 mL). The reaction solution was warmed to room temperature and, after an additional 1 h, methyl-pyridin-3-ylmethyl-amine (180 mg) was added. After an additional 1 h, the solution was concentrated, the resulting residue was dissolved in EtOAc, and the solution washed sequentially with saturated aqueous NH_4Cl solution and water, dried over Na_2SO_4 , and concentrated to afford the product carbamate as a mixture of isomers that was used without further purification.

Step E

A solution of the product of Step D (230 mg) in CH_3CN was heated to 150°C in a microwave apparatus for 5 min. The solution was concentrated and the resulting residue was purified by silica gel chromatography to afford the title compound as a white solid. $\text{MS (M-H)}^- = 384.3$.

The following compounds were prepared in a manner analogous to that described in Example 221 using appropriate starting materials.

Example	Name	MS (M+H) ⁺ or MS (M-H) ⁻
222	<i>tert</i> -Butyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	337.1 (+)

223	(6-Methyl-pyridin-2-ylmethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	386.1 (+)
224	(3-Chloro-5-trifluoromethyl-pyridin-2-ylmethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	474 (+)
225	(1,1-Dimethyl-propyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	351.3 (+)
226	[2-(4-Fluoro-phenyl)-1,1-dimethyl-ethyl]-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	431.3 (+)
227	(1-Methyl-1-phenyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	399.3 (+)
228	Diethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	337.3 (+)
229	(2,2,2-Trifluoro-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	363.2 (+)
230	<i>tert</i> -Butyl-methyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	351.3 (+)
231	(Tetrahydro-pyran-2-ylmethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	379.3 (+)
232	Methyl-propyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	337.2 (+)
233	Isobutyl-methyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	351.2 (+)
234	Cyclopropylmethyl-propyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	377.3 (+)
235	(2-Methoxy-ethyl)-methyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	353.2 (+)
236	Cyclohexyl-methyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-	377.3 (+)

	2H-pyrazol-3-yl]-cyclobutyl ester	
237	2-Ethyl-piperidine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	377.3 (+)
238	(2-Cyano-ethyl)-cyclohexyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	416.2 (+)
239	Ethyl-isopropyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	351.2 (+)
240	(R)-2-Methoxymethyl-pyrrolidine-1-carboxylic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	379.2 (+)
241	Methyl-((R)-1-phenyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	399.2 (+)
242	Isopropyl-(2-methoxy-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	381.2 (+)
243	Methyl-(3-methyl-pyridin-2-ylmethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	398.3(-)
244	Methyl-quinoxalin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	437.3 (+)
245	Methyl-quinolin-8-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	436.3 (+)
246	Methyl-quinolin-6-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	436.2 (+)
247	Methyl-quinolin-5-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	436.2 (+)
248	Methyl-pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	386.2 (+)
249	Methyl-pyridin-4-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-	386.2 (+)

	propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	
250	Methyl-(1-pyridin-2-yl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	400.1 (+)
251	(1-Phenyl-cyclopentyl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	425.2 (+)
252	Isobutyl-carbamic acid <i>cis</i> -3-{5-[(2-methyl-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	365.2 (+)
253	Isobutyl-carbamic acid <i>cis</i> -3-{5-[(4-methyl-tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	379.4 (+)
254	<i>tert</i> -Butyl-carbamic acid <i>cis</i> -3-{5-[(2-methyl-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	365.3 (+)
255	Dimethyl-carbamic acid <i>cis</i> -3-{5-[(2-methyl-tetrahydro-furan-2-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	337.2 (+)
256	<i>tert</i> -Butyl-carbamic acid <i>cis</i> -3-{5-[(4-methyl-tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	379.4 (+)
257	Dimethyl-carbamic acid <i>cis</i> -3-{5-[(4-methyl-tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	351.4 (+)
258	Benzyl-carbamic acid <i>cis</i> -3-{5-[(4-methyl-tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	413.4 (+)
259	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-{5-[(4-methyl-tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	414.5 (+)

The following compounds were prepared in a manner analogous to that described in Example 221 using appropriate starting materials, except Step E employed DMSO as solvent, rather than CH₃CN.

Example	Name	MS (M+H) ⁺
---------	------	--------------------------

260	Propyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	386.3
261	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	435.2
262	Isopropyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	386.4
263	Cyclobutyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	398.3
264	Methyl-propyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	400.3
265	((S)-1-Phenyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	448.3
266	((R)-1-Phenyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	448.3
267	Isopropyl-methyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	400.3
268	Methyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	358.2
269	(2-Methoxy-ethyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	402.3
270	Diethyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	400.3
271	<i>tert</i> -Butyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	400.3
272	(1-Methyl-1-phenyl-ethyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-	462.4

	yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	
273	Ethyl-methyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	386.3

Example 274**Cyclopropylmethyl-carbamic acid 3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester**

- 5 A mixture of 5-(3-cyclopropylmethylcarbamoyloxy-cyclobutyl)-3-(2-methyl-2-pyridin-2-yl-propionylamino)-pyrazole-1-carboxylic acid *tert*-butyl ester and 3-(3-cyclopropylmethylcarbamoyloxy-cyclobutyl)-5-(2-methyl-2-pyridin-2-yl-propionylamino)-pyrazole-1-carboxylic acid *tert*-butyl ester (120 mg, prepared by procedures analogous to those described in Example 221, Steps A to E, using appropriate starting materials) was dissolved in TFA (0.19 mL) and CH₂Cl₂ (1.25 mL) at room temperature. After 4 h, the solution was diluted with EtOAc and the solution was washed sequentially with saturated aqueous NaHCO₃ solution (2x), water, and saturated aqueous NaCl solution. The organic layer was dried over Na₂SO₄ and concentrated. Purification of the residue by silica gel chromatography afforded the title compound as a solid. MS (M+H)⁺ = 398.3.

The following compounds were prepared in a manner analogous to that described in Example 274 using appropriate starting materials.

Example	Name	MS (M+H) ⁺
275	<i>tert</i> -Butyl-carbamic acid <i>cis</i> -3-[5-(cyclohexanecarbonyl-amino)-2H-pyrazol-3-yl]-cyclobutyl ester	363.5
276	Phenyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	357.3
277	Methyl-pyridin-2-yl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	372.3
278	(6-Methyl-pyridin-2-yl)-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	372.3
279	Methyl-phenyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	371.3
	Pyrimidin-2-yl-carbamic acid <i>cis</i> -3-[5-	

280	(2,2-dimethyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	359.3
281	Ethyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	372.3
282	(2-Fluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	452.2
283	Pyrrolidine-1-carboxylic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	398.3
284	Morpholine-4-carboxylic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	414.3
285	(2,6-Difluoro-benzyl)-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	470.2
286	Methyl-pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	449.0
287	<i>tert</i> -Butyl-methyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	414.4

Example 2882-Methyl-pyrrolidine-1-carboxylic acid *cis*-3-[5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester5 Step A

A mixture of 5-(*cis*-3-hydroxy-cyclobutyl)-3-[(tetrahydro-pyran-4-carbonyl)-amino]-pyrazole-1-carboxylic acid *tert*-butyl ester and 3-(*cis*-3-hydroxy-cyclobutyl)-5-[(tetrahydro-pyran-4-carbonyl)-amino]-pyrazole-1-carboxylic acid *tert*-butyl ester (120 mg, prepared by procedures analogous to those described in Example 221, Steps B to C, using appropriate starting materials), triphosgene (69 mg), and PS-DMAP resin (0.30 mmol) were stirred in CH₂Cl₂ (2 mL) at 0 °C. After 60 min 2-methyl-pyrrolidine (0.10 mL) was added and the resulting mixture was stirred overnight. The resin was removed by filtration and the filtrate was concentrated to afford the product carbamate that was used without further purification.

Step B

A solution of the product of Step A (157 mg) in DMSO was heated at 150 °C in a microwave apparatus for 7 min. The solution was then purified by reversed-phase preparative HPLC to afford the title product as a solid. MS (M+H)⁺ = 377.2.

- 5 The following compounds were prepared in a manner analogous to that described in Example 288 using appropriate starting materials.

Example	Name	MS (M+H) ⁺ or MS (M-H) ⁻
289	<i>tert</i> -Butyl-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	363.2 (-)
290	(6-Methyl-pyridin-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	412.3 (-)
291	(1-Methyl-1-phenyl-ethyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	427.2 (+)
292	Methyl-(3-methyl-pyridin-2-ylmethyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	426.1 (-)
293	(2,2,2-Trifluoro-1-pyridin-2-yl-ethyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	468.1 (+)
294	(2R,6S)-2,6-Dimethyl-morpholine-4-carboxylic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	407.2 (+)
295	2-Pyridin-2-yl-pyrrolidine-1-carboxylic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	438.1 (-)
296	Methyl-pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	414.2 (+)

297	Methyl-(1-pyridin-2-yl-ethyl)-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	426.1 (-)
298	2-Methoxymethyl-pyrrolidine-1-carboxylic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	407.2 (+)
299	<i>tert</i> -Butyl-methyl-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	379.2 (+)
300	Benzyl-ethyl-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-pyran-4-carbonyl)-amino]-2H-pyrazol-3-yl}-cyclobutyl ester	427.2 (+)

Example 301

Isobutyl-carbamic acid *cis*-3-[5-(2-*p*-tolyl-acetyl-amino)-1H-pyrazol-3-yl]-cyclobutyl ester

5 Step A

A solution of 2-*tert*-butyl-5-(3,3-dimethoxy-cyclobutyl)-2H-pyrazol-3-ylamine (4.0 g), di-*tert*-butyl dicarbonate (10.4 g), Et₃N (6.6 mL), and DMAP (40 mg) in CH₂Cl₂ (60 mL) was stirred at room temperature for 3 days. The solution was concentrated to afford the *bis*-carbamoyleated product as an oil that was used without further purification.

10 Step B

The product of Step A was converted into [2-*tert*-butyl-5-(*cis*-3-isobutylcarbamoyleoxy-cyclobutyl)-2H-pyrazol-3-yl]-imidodicarboxylic acid di-*tert*-butyl ester using appropriate starting materials by procedures analogous to those described in Example 1, Steps B to C, and Example 9, Step A.

15 Step C

A solution of the product of Step B (4.5 g) in TFA (40 mL) was stirred at room temperature for 4 h. The solution was concentrated and the residue was dissolved in CHCl₃. The solution was washed sequentially with saturated aqueous NaHCO₃ solution and saturated aqueous NaCl solution. The organic layer was dried over Na₂SO₄ and concentrated. Purification of the residue by silica gel chromatography afforded the amine product as an oil.

Step D

To *p*-tolyl-acetic acid (0.18 mmol) was added sequentially: the product of Step C (1 mL of a 0.16 M solution in EtOAc), 1-propanephosphonic acid cyclic anhydride (1 mL of a 0.36 M solution in EtOAc), and Et₃N (1 mL of a 0.72 M solution in EtOAc).

- 5 The resulting solution was heated at reflux overnight and concentrated to afford a residue that was used without purification in the next step.

Step E

- The product of Step D was dissolved in TFA (2 mL) and the solution was heated at reflux overnight. The solution was concentrated and the residue purified by reversed-phase preparative HPLC to afford the title product. MS (M+H)⁺ = 385.4.

The following compounds were prepared in a manner analogous to that described in Example 301 using appropriate starting materials.

Example	Name	MS (M+H) ⁺
302	Isobutyl-carbamic acid <i>cis</i> -3-[5-(3-methyl-2-phenyl-butyrylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	413.5
303	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2- <i>m</i> -tolyl-acetylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	385.4
304	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(3-methoxy-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	401.4
305	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(4-chloro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	405.4
306	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(4-methoxy-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	401.4
307	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2- <i>o</i> -tolyl-acetylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	385.4
308	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(3-fluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	389.4
309	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(3,4-dimethoxy-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	431.4
310	Isobutyl-carbamic acid <i>cis</i> -3-[5-[1-(4-chloro-phenyl)-cyclopentanecarbonyl]-amino]-1H-pyrazol-3-yl]-cyclobutyl ester	459.4
311	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(2-methoxy-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	401.4

312	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(3-trifluoromethyl-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	439.4
313	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(2-trifluoromethyl-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	439.4
314	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(2,6-dichloro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	439.3
315	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(4-isopropyl-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	413.4
316	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(2-chloro-4-fluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	423.3
317	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(2,4-difluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	407.4
318	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(2,6-difluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	407.4
319	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(3,4-dichloro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	439.3
320	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(2,5-difluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	407.4
321	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(3,4-difluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	407.4
322	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2-benzo[1,3]dioxol-5-yl-acetylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	415.4
323	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(4-fluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	389.4
324	Isobutyl-carbamic acid <i>cis</i> -3-(5-phenylacetylamino-1H-pyrazol-3-yl)-cyclobutyl ester	371.4
325	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(4-chloro-phenyl)-2-methyl-propionylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	433.4
326	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2,3-diphenyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	461.5
327	Isobutyl-carbamic acid <i>cis</i> -3-[5-[2-(3,5-bis-trifluoromethyl-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	507.5

328	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(2,4,5-trifluoro-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	425.4
329	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(4-methoxy-3-methyl-phenyl)-acetylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	415.5
330	Isobutyl-carbamic acid <i>cis</i> -3-(5-{[1-(4-fluoro-phenyl)-cyclopentanecarbonyl]-amino}-1H-pyrazol-3-yl)-cyclobutyl ester	443.5
331	Isobutyl-carbamic acid <i>cis</i> -3-(5-phenylacetylamino-2H-pyrazol-3-yl)-cyclobutyl ester	371.4
332	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(3-chloro-phenyl)-acetylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	405.1
333	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(4-bromo-phenyl)-acetylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	449.1
334	Isobutyl-carbamic acid <i>cis</i> -3-[5-(3-methyl-2-phenyl-pentanoylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	427.5
335	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(2,4-dichloro-phenyl)-acetylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	439.3
336	Isobutyl-carbamic acid <i>cis</i> -3-(5-{[1-(4-chloro-phenyl)-cyclopropanecarbonyl]-amino}-2H-pyrazol-3-yl)-cyclobutyl ester	431.3
337	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(3,5-difluoro-phenyl)-acetylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	407.4
338	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2-benzo[1,3]dioxol-5-yl-acetylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	415.5
339	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2- <i>p</i> -tolyl-acetylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	385.4
340	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(2-bromo-5-chloro-phenyl)-acetylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	485.2
341	Isobutyl-carbamic acid <i>cis</i> -3-[5-(2-isochroman-7-yl-acetylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	427.5
342	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(2-chloro-6-fluoro-phenyl)-acetylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	423.9
343	Isobutyl-carbamic acid <i>cis</i> -3-{5-[2-(3-acetyl-phenyl)-acetylamino]-2H-	413.3

	pyrazol-3-yl]-cyclobutyl ester	
--	--------------------------------	--

Example 344**Cyclobutyl-carbamic acid *cis*-3-[5-(3-pyridin-3-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester****5 Step A**

A solution of cyclobutyl-carbamic acid *cis*-3-(5-amino-1-*tert*-butyl-1H-pyrazol-3-yl)-cyclobutyl ester (100 mg, prepared by procedures analogous to those described in Example 301, Steps A to C, using appropriate starting materials), 1-propanephosphonic acid cyclic anhydride (50 wt. % in EtOAc, 407 mg), and 3-pyridin-3-yl-propionic acid (60 mg) in EtOAc (2 mL) was heated at reflux overnight. The mixture was concentrated and the residue was used without purification in the next step.

Step B

A solution of the product of Step A in TFA (4 mL) was heated at 80 °C overnight. The solution was then concentrated and the residue was purified by reversed-phase preparative HPLC to afford the title product. MS (M+H)⁺ = 384.4.

The following compounds were prepared in a manner analogous to that described in Example 344 using appropriate starting materials. For Examples 345 to 350 and Examples 352 to 368, Et₃SiH (3 equivalents) was added to the reaction mixture in Step B.

Example	Name	MS (M+H) ⁺
345	Cyclopentyl-carbamic acid <i>cis</i> -3-[5-[3-(4-methyl-thiazol-5-yl)-propionylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	418.4
346	Cyclopentyl-carbamic acid <i>cis</i> -3-[5-[3-(1-methyl-1H-pyrazol-4-yl)-propionylamino]-1H-pyrazol-3-yl]-cyclobutyl ester	401.5
347	Cyclopentyl-carbamic acid <i>cis</i> -3-[5-(3-pyridin-3-yl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	398.5
348	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester	372.5
349	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-[2-(4-chloro-phenoxy)-2-methyl-propionylamino]-1H-pyrazol-	484.5

	3-yl]-cyclobutyl ester	
350	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-{5-[2-(2-chloro-phenoxy)- acetylamino]-1H-pyrazol-3-yl]- cyclobutyl ester	456.4
351	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-(5-isobutyrylamino-1H-pyrazol- 3-yl)-cyclobutyl ester	358.4
352	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(3,3-dimethyl-butyrylamino)- 1H-pyrazol-3-yl]-cyclobutyl ester	386.5
353	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(3-methyl-butyrylamino)-1H- pyrazol-3-yl]-cyclobutyl ester	372.5
354	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(cyclopentanecarbonyl- amino)-1H-pyrazol-3-yl]-cyclobutyl ester	384.5
355	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(cycloheptanecarbonyl- amino)-1H-pyrazol-3-yl]-cyclobutyl ester	412.5
356	Methyl-carbamic acid <i>cis</i> -3-[5-(3- phenyl-propionylamino)-1H-pyrazol- 3-yl]-cyclobutyl ester	343.4
357	Dimethyl-carbamic acid <i>cis</i> -3-[5-(3- phenyl-propionylamino)-1H-pyrazol- 3-yl]-cyclobutyl ester	357.4
358	Methyl-carbamic acid <i>cis</i> -3-[5-[2-(2- chloro-phenoxy)-acetylamino]-1H- pyrazol-3-yl]-cyclobutyl ester	379.4
359	Dimethyl-carbamic acid <i>cis</i> -3-[5-[2- (2-chloro-phenoxy)-acetylamino]- 1H-pyrazol-3-yl]-cyclobutyl ester	393.4
360	Methyl-carbamic acid <i>cis</i> -3-[5-(2- phenoxy-acetylamino)-1H-pyrazol-3- yl]-cyclobutyl ester	345.3
361	Dimethyl-carbamic acid <i>cis</i> -3-[5-(2- phenoxy-acetylamino)-1H-pyrazol-3- yl]-cyclobutyl ester	359.4
362	Dimethyl-carbamic acid <i>cis</i> -3-[5- (2,2-dimethyl-propionylamino)-1H- pyrazol-3-yl]-cyclobutyl ester	309.4
363	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-[(pyridine-2-carbonyl)- amino]-1H-pyrazol-3-yl]-cyclobutyl ester	393.4
364	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-[(<i>S</i>)-tetrahydro-furan-2- carbonyl]-amino]-1H-pyrazol-3-yl]-	386.2

	cyclobutyl ester	
365	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-{5-[2-(4-chloro-phenyl)-2-methyl-propionylamino]-1H-pyrazol-3-yl}-cyclobutyl ester	468.2
366	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-{5-[(5-methyl-pyrazine-2-carbonyl)-amino]-1H-pyrazol-3-yl}-cyclobutyl ester	408.3
367	Pyridin-2-ylmethyl-carbamic acid <i>cis</i> -3-[5-(2-pyridin-2-yl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	421.1
368	Dimethyl-carbamic acid <i>cis</i> -3-[5-(3-cyclohexyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	363.2
369	Cyclobutyl-carbamic acid <i>cis</i> -3-{5-[(tetrahydro-furan-2-carbonyl)-amino]-1H-pyrazol-3-yl}-cyclobutyl ester	349.4

Example 370Cyclobutyl-carbamic acid *cis*-3-[5-(2,2-dimethyl-propionylamino)-1H-pyrazol-3-yl]-cyclobutyl ester5 Step A

A solution of cyclobutyl-carbamic acid *cis*-3-(5-amino-1-*tert*-butyl-1H-pyrazol-3-yl)-cyclobutyl ester (100 mg, prepared by procedures analogous to those described in Example 301, Steps A to C, using appropriate starting materials), trimethylacetyl chloride (46 mg), and Et₃N (50 mg) in CH₂Cl₂ was stirred overnight at room temperature. The solution was concentrated to afford a residue that was used without further purification.

10 Step B

A solution of the product of Step A in TFA (4 mL) was heated at 80 °C overnight. The solution was concentrated and the residue purified by reversed-phase preparative HPLC to afford the title product. MS (M+H)⁺ = 335.4

The following compounds were prepared in a manner analogous to that described in Example 370 using appropriate starting materials. For Examples 373 to 377, Et₃SiH (3 equivalents) was added to the reaction mixture in Step B.

Example	Name	MS (M+H) ⁺
371	Methyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-propionylamino)-1H-	295.4

	pyrazol-3-yl]-cyclobutyl ester	
372	Dimethyl-carbamic acid <i>cis</i> -3-{5-[(1-phenyl-cyclopentanecarbonyl)-amino]-2H-pyrazol-3-yl]-cyclobutyl ester	397.2
373	Dimethyl-carbamic acid <i>cis</i> -3-[5-(2-methyl-2-phenyl-propionylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	371.2
374	Dimethyl-carbamic acid <i>cis</i> -3-(5-[(1-(4-chloro-phenyl)-cyclobutanecarbonyl]-amino)-2H-pyrazol-3-yl)-cyclobutyl ester	417.2
375	Dimethyl-carbamic acid <i>cis</i> -3-{5-[2-(2-chloro-phenoxy)-2-methyl-propionylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	421.1
376	Dimethyl-carbamic acid <i>cis</i> -3-{5-[2-(4-chloro-phenoxy)-2-methyl-propionylamino]-2H-pyrazol-3-yl]-cyclobutyl ester	421.1
377	Dimethyl-carbamic acid <i>cis</i> -3-[5-(2,2-dimethyl-pentanoylamino)-2H-pyrazol-3-yl]-cyclobutyl ester	337.2